

GLOBAL EXISTENCE FOR A COUPLED WAVE SYSTEM RELATED TO THE STRAUSS CONJECTURE

JASON METCALFE AND DAVID SPENCER

ABSTRACT. A coupled system of semilinear wave equations is considered, and a small data global existence result related to the Strauss conjecture is proved. Previous results have shown that one of the powers may be reduced below the critical power for the Strauss conjecture provided the other power sufficiently exceeds such. The stability of such results under asymptotically flat perturbations of the space-time where an integrated local energy decay estimate is available is established.

1. INTRODUCTION

The purpose of this article is to establish global existence for a coupled system of wave equations, which is related to the Strauss conjecture, on asymptotically flat space-times that permit a localized energy estimate. It is now well-known that nonlinear wave equations

$$\square u := (\partial_t^2 - \Delta)u = F_p(u), \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^n$$

where

$$(1.1) \quad \sum_{0 \leq j \leq 2} |u|^j |\partial_u^j F_p(u)| \lesssim |u|^p \quad \text{for } u \text{ small}$$

have global solutions for sufficiently small initial data provided $p > p_c$ where $p_c > 1$ solves

$$(1.2) \quad (n-1)p_c^2 - (n+1)p_c - 2 = 0.$$

Moreover, blow up is known to occur for $p < p_c$. These results originated in [23] for $n = 3$ where $p_c = 1 + \sqrt{2}$, and following [44] the problem became known as the Strauss conjecture. Global existence in general dimension was eventually established in [20], [45]; see the references therein for many intermediate results. Blow up below the critical exponent was proved in [40]. See also [39], [51] for further results at the critical exponent.

In the current work, we shall examine a system of the form

$$\square u = |v|^p, \quad \square v = |u|^q.$$

In the flat case, the coupled system was examined in [16], and it was shown that global existence may be established for powers in the nonlinearity below the critical

Date: January 23, 2017.

The first author was supported in part by NSF grant DMS-1054289. The second author was supported in part by a Summer Undergraduate Research Fellowship (SURF) through the University of North Carolina, and the results contained herein were developed as a part of his Undergraduate Honors Thesis.

exponent provided the power on the coupled equation exceeds the same. Indeed, setting

$$(1.3) \quad C(p, q) = \max \left\{ \frac{q + 2 + p^{-1}}{pq - 1}, \frac{p + 2 + q^{-1}}{pq - 1} \right\} - \frac{n - 1}{2},$$

it was shown that small data global existence holds for $C(p, q) < 0$ and that such fails for $C(p, q) > 0$. Notice that $C(p, p) = 0$ corresponds precisely to (1.2). In particular, note that small data global existence may be established for powers $p < p_c$ provided that the other power q sufficiently exceeds p_c . In addition to [16], see [17], [14] for treatments of $C(p, q) > 0$ and [1], [15], [25], and [21] for analysis of the critical curve $C(p, q) = 0$. Moreover, see the overview [26] of this and related problems.

Here we seek to establish the same using techniques that are sufficiently robust so as to allow background geometries. Specifically, we shall use a variant of the weighted Strichartz estimates of [22], [19], which were further developed in [27], [35], and the localized energy estimate to prove such global existence.

We shall examine operators of the form

$$(1.4) \quad Pu = \partial_\alpha g^{\alpha\beta} \partial_\beta u + b^\alpha \partial_\alpha u + cu$$

on space-times M where $M = \mathbb{R}_+ \times \mathbb{R}^3$ or $M = \mathbb{R}_+ \times (\mathbb{R}^3 \setminus \mathcal{K})$ where \mathcal{K} has a smooth boundary and $\mathcal{K} \subset \{x : |x| < R_0\}$. Here g is a Lorentzian metric, and we make the assumption that g can be written as

$$(1.5) \quad g(t, x) = m + g_0(t, r) + g_1(t, x)$$

where $m = \text{diag}(-1, 1, 1, 1)$ is the Minkowski metric. The components g_0 and g_1 will represent long-range and short-range perturbations respectively. They are asymptotically flat in the sense that

$$(1.6) \quad \|\partial_{t,x}^\mu g_{i,\alpha\beta}\|_{\ell_1^{i+|\mu|} L_{t,x}^\infty} = \mathcal{O}(1), \quad i = 0, 1, \quad |\mu| \leq 3.^1$$

Due to the need to commute with spatial rotations, the long-range perturbation g_0 is assumed to be spherically symmetric in the sense that the coefficients only (spatially) depend on $r = |x|$ and

$$(1.7) \quad g - g_1 = (-1 + \tilde{g}_{00}(t, r)) dt^2 + 2\tilde{g}_{01}(t, r) dt dr + (1 + \tilde{g}_{11}(t, r)) dr^2 + (1 + \tilde{g}_{22}(t, r)) r^2 d\omega_{\mathbb{S}^2}^2.$$

and, by (1.6), $\|\partial_{t,x}^\mu \tilde{g}_{\alpha\beta}\|_{\ell_1^{|\mu|} L_{t,x}^\infty} = \mathcal{O}(1)$ for $|\mu| \leq 3$. The coefficients of the lower-order perturbations decay are assumed to decay as follows:

$$(1.8) \quad \|\partial_{t,x}^\mu b\|_{\ell_1^{1+|\mu|} L_{t,x}^\infty} + \|\partial_{t,x}^\mu c\|_{\ell_1^{2+|\mu|} L_{t,x}^\infty} = \mathcal{O}(1), \quad |\mu| \leq 2.$$

We shall also assume that the perturbations admit a (weak) localized energy decay. More specifically, we assume that there is R_1 (with $R_1 > R_0$ in the case

¹Here, as in [35], for a norm A , we set

$$\|u\|_{\ell_q^s A} = \left\| 2^{js} \|\phi_j(x) u(t, x)\|_A \right\|_{\ell_{j \geq 0}^q}, \quad \sum_{j \geq 0} \phi_j^2(x) = 1, \quad \text{supp } \phi_j \subset \{\langle x \rangle \approx 2^j\}.$$

that $M = \mathbb{R}_+ \times (\mathbb{R}^3 \setminus \mathcal{K})$ so that if u solves $Pu = F$ then

$$(1.9) \quad \|\partial \partial^\mu u\|_{L_t^\infty L_x^2} + \|(1 - \chi) \partial \partial^\mu u\|_{\ell_\infty^{-1/2} L_{t,x}^2} + \|\partial^\mu u\|_{\ell_\infty^{-3/2} L_{t,x}^2} \\ \lesssim \|u(0, \cdot)\|_{H^{|\mu|+1}} + \|\partial_t u(0, \cdot)\|_{H^{|\mu|}} + \sum_{|\nu| \leq |\mu|} \|\partial^\nu F\|_{L_t^1 L_x^2}$$

for all $|\mu| \leq 2$. Here χ is a smooth function that is identically 1 on $B_{R_1/2} := \{|x| \leq R_1/2\}$ and is supported on B_{R_1} .

On $(1+3)$ -dimensional Minkowski space (i.e. when $g_0 \equiv g_1 \equiv 0$), it is known that

$$\|\partial u\|_{L_t^\infty L_x^2} + \|\partial u\|_{\ell_\infty^{-1/2} L_{t,x}^2} + \|u\|_{\ell_\infty^{-3/2} L_{t,x}^2} \lesssim \|\partial u(0, \cdot)\|_{L^2} + \|\square u\|_{L_t^1 L_x^2},$$

which is a stronger version of the $\mu = 0$ estimate above. And as the flat d'Alembertian commutes with $\partial_{t,x}$, the higher order variants readily follow. Such estimates originated in [36]. They follow, e.g., by multiplying $\square u$ by a multiplier of the form $\frac{r}{r+2\tau} \partial_r u + \frac{n-1}{2} \frac{1}{r+2\tau} u$, integrating over $[0, T] \times \mathbb{R}^3$, and integrating by parts. See, e.g., [43], [29]. And see, e.g., [34] for a more complete history. These estimates are known to be rather robust in the asymptotically flat regime. Even without the cutoff, they are known to hold for small, possibly time-dependent perturbations of Minkowski space [29, 30], [33, 32], [2] and for time-independent nontrapping perturbations in the product manifold setting due to, e.g., [8], [6], [42]. See [31] for the most general results in the nontrapping regime.

The presence of trapped rays is a known obstruction to the localized energy estimate [37], [38]. The asymptotic flatness restricts the possibility of trapped rays to a compact set, and when the trapping is sufficiently weak, a localized energy estimate where one, say, cuts off away from the trapping may sometimes be recovered. Allowing for this is the reason for the cutoff in assumption (1.9). Previous results have then verified (1.9) in a number of settings where trapping occurs, including on the Schwarzschild space-time [4, 5], [10, 11], [28], on Kerr space-times with $a \ll M$ [47] (see also [3], [9, 12] for some closely related results and [13] for a related result that holds for the full subextremal range $|a| < M$), and on certain warped product manifolds that contain degenerate trapping [7].

We now introduce the specific problem at hand. With two possibly different operators P_1 and P_2 subject to hypotheses (1.5)-(1.9), we examine the coupled system


$$(1.10) \quad \begin{aligned} P_1 u &= F_p(v), & P_2 v &= F_q(u), \\ u(0, x) &= f_1(x), & v(0, x) &= f_2(x), \\ \partial_t u(0, x) &= g_1(x), & \partial_t v(0, x) &= g_2(x). \end{aligned}$$

Here F_p and F_q are two functions satisfying (1.1).

For the system (1.10), we shall establish the following small data global existence result:

Theorem 1.1. *Suppose that P_1 and P_2 are operators of the form (1.4) so that (1.5), (1.6), (1.7), (1.8), and (1.9) hold. Moreover assume that $2 < p, q$ and $C(p, q) < 0$.*

FIGURE 1. Range of allowable indices



Then if $f_1, g_1, f_2, g_2 \in C_c^\infty$ ² and

$$(1.11) \quad \|(f_1, f_2)\|_{H^3} + \|(g_1, g_2)\|_{H^2} \leq \varepsilon$$

with ε sufficiently small, there exists a global solution (u, v) to (1.10).

We note that the techniques to prove Theorem 1.1 also work in four spatial dimensions, but as they require $p, q \geq 2$ and $p_c = 2$ in this case, nothing new is gained over [35].

This result follows a number of studies that established various existence results related to the Strauss conjecture in the presence of background geometry. In exterior domains, these included [18], [22], [41], and [52]. And on asymptotically flat backgrounds, see [42], [49], [27], [35], and [48]. See, also, the expository article [50]. A key component of many of these results is the weighted Strichartz estimate of [22], [19]. Here, in particular, we rely on the variant of that developed in [35], which is based on the local energy estimates of [33].

Figure 1 demonstrates the range of allowable indices. Theorem 1.1 allows for any pair of indices (p, q) that land outside of the shaded region in the lower left corner. The techniques of, e.g., [35], however, only trivially apply in the rectangular region where $p, q > p_c = 1 + \sqrt{2}$. The other curve will be discussed further in Section 3.

The method that we shall employ is similar in spirit to that of [35], which in turn is based on a number of preceding works. Near infinity, where the asymptotic flatness allows us to think of the geometry as a small perturbation of Minkowski space, variants of the weighted Strichartz estimates of [22], [19] are employed. The assumed localized energy estimate handles the remaining compact region, where the geometry has the most significant role. It also allows the analyses done in the two regions to be glued together. Such a strategy has become common. See, e.g., [27], [28], [33, 32], [34], [46].

²For simplicity of exposition, we have taken the data here to be compactly supported, but this may be replaced by a condition such as [35, (5.3)].

2. MAIN ESTIMATES

Before we proceed to the main estimates for the linear equation, we first introduce some notation. We shall be using a restricted set of the classical invariant vector fields. We let $\Omega_{ij} = x_i \partial_j - x_j \partial_i$ denote the generators of spatial rotations. And we let $Y = \{\nabla_x, \Omega\}$ and $Z = \{\partial, \Omega\}$, where $\partial = (\partial_t, \nabla_x)$ denotes the space-time gradient. We also introduce the shorthand $|Z^{\leq m} u| = \sum_{|\mu| \leq m} |Z^\mu u|$ for summing over multi-indices of order $\leq m$. A similar notation where the absolute values are replaced by a norm shall also be used. Finally, for the mixed norms that appear in the weighted Strichartz estimates below, we fix the convention

$$\|f\|_{L_t^p L_r^q L_\omega^s} = \left[\int \left(\int \left[\int |f(t, r\omega)|^s d\omega_{\mathbb{S}^2} \right]^{q/s} r^2 dr \right)^{p/q} dt \right]^{1/p},$$

with the obvious changes for Lebesgue indices of ∞ .

The main linear estimate that will be applied near infinity, where the operators may be viewed as small perturbations of the flat d'Alembertian, is the following weighted Strichartz estimate:

Theorem 2.1 ([35]). *Suppose that P is an operator of the form (1.4) satisfying the hypotheses (1.5), (1.6), (1.7), (1.8), and (1.9). Suppose that $w(0, \cdot)$ and $\partial_t w(0, \cdot)$ are compactly supported and satisfy (1.11). Then there exists $R_2 > R_1$ so that for any $R > R_2$ if ψ_R is identically 1 on $\{|x| \geq 2R\}$ and vanishes on $\{|x| < R\}$, then we have*

$$(2.1) \quad \|\psi_R Z^{\leq 2} w\|_{\ell_p^{\frac{3}{2} - \frac{4}{p} - s} L_t^p L_r^p L_\omega^2} \leq C_1 \varepsilon + C_1 \|\psi_R Y^{\leq 1} Pw(0, \cdot)\|_{\dot{H}^{s-1}} \\ + C_1 \|\psi_R^{\tilde{p}} Z^{\leq 2} Pw\|_{\ell_1^{-\frac{1}{2} - s} L_t^1 L_r^1 L_\omega^2} + C_1 \|\partial^{\leq 2} Pw\|_{L_t^1 L_r^2 L_\omega^2}$$

and

$$(2.2) \quad \|\psi_R w\|_{\ell_p^{\frac{3}{2} - \frac{4}{p} - s} L_t^p L_r^p L_\omega^2} \lesssim \varepsilon + \|\psi_R^{\tilde{p}} Pw\|_{\ell_1^{-\frac{1}{2} - s} L_t^1 L_r^1 L_\omega^2} + \|Pw\|_{L_t^1 L_r^2 L_\omega^2}$$

for any $p \in (2, \infty)$, $s \in (1/2 - 1/p, 1/2)$, and $\tilde{p} > 0$.

These estimates originated in [22] and [19] for the flat d'Alembertian. The above version is essentially from [35], which in particular allows for asymptotically flat operators and does not necessitate compactly supported initial data though we assume that here for simplicity. These estimates follow by interpolating a variant of the localized energy estimate with a trace theorem on the sphere. We note a few minor modifications from the version in [35]: (1) We have allowed a different power of the cutoff function in the right sides; (2) We have more carefully stated the requirements on R so that in the sequel we are able to use the same R for both u and v ; (3) We have stated separately the case where no vector fields are applied since this is used when showing that our iteration is Cauchy. These all follow from trivial modifications of the proof of [35].

A key to tying the region near infinity to the remaining compact region where (1.9) is the primary tool is the following weighted Sobolev inequalities. These are variants of the original estimates of [24] and follow by localizing, applying Sobolev embeddings on $\mathbb{R} \times \mathbb{S}^2$, and adjusting the volume elements to match those of \mathbb{R}^3 in polar coordinates. See, e.g., [27], [35] for proofs.

Lemma 2.2. *On \mathbb{R}^3 , for $R \geq 1$, $\beta \in \mathbb{R}$, and $2 \leq p \leq q \leq \infty$, we have*

$$(2.3) \quad \|r^\beta u\|_{L_r^q L_\omega^\infty(r \geq R+1)} \lesssim \sum_{|\mu| \leq 2} \|r^{\beta - \frac{2}{p} + \frac{2}{q}} Y^\mu u\|_{L_r^p L_\omega^2(r \geq R)}.$$

If $2 \leq p \leq q \leq 4$ and $\beta \in \mathbb{R}$, we also have

$$(2.4) \quad \|r^\beta u\|_{L_r^q L_\omega^4(r \geq R+1)} \lesssim \sum_{|\mu| \leq 1} \|r^{\beta - \frac{2}{p} + \frac{2}{q}} Y^\mu u\|_{L_r^p L_\omega^2(r \geq R)}.$$

3. SMALL DATA GLOBAL EXISTENCE

We now prove Theorem 1.1. We shall apply (2.1) to u with $(p, s) = \left(q, \frac{7+4p-3pq}{2-2pq}\right)$ and to v with $(p, s) = \left(p, \frac{7+4q-3pq}{2-2pq}\right)$. The requirement from Theorem 2.1 that $s > \frac{1}{2} - \frac{1}{p}$ then corresponds to

$$\frac{2+p+q^{-1}}{pq-1} < 1, \quad \frac{2+q+p^{-1}}{pq-1} < 1$$

respectively. In the case of $n = 3$, this produces the requirement that $C(p, q) < 0$.

We shall assume, without loss of generality, that (p, q) satisfy

$$(3.1) \quad p(q-2) < 3, \quad q(p-2) < 3,$$

which corresponds to the condition that $s < 1/2$ in Theorem 2.1. It is these conditions that are represented by the curve in Figure 1, below which they are satisfied. In the unshaded region to the left of the rectangle, if (p, q) satisfy $C(p, q) < 0$ but one of the above conditions is violated, one may simply choose any $\tilde{q} < q$ so that $C(p, \tilde{q}) < 0$ and so that both conditions in (3.1) hold. One simply imagines $|u|^q = |u|^{q-\tilde{q}}|u|^{\tilde{q}}$ and argues as below with the exponent \tilde{q} replacing q . Simple Sobolev embeddings control the remaining $q - \tilde{q}$ powers. In the unshaded region below the rectangle, one argues similarly by reducing the power of p . In the shaded rectangle, the methods of [35] apply directly and no further argument is needed.

Let $s_1 = \frac{7+4p-3pq}{2-2pq}$, $s_2 = \frac{7+4q-3pq}{2-2pq}$, $\alpha_1 = \frac{3}{2} - \frac{4}{q} - s_1$, and $\alpha_2 = \frac{3}{2} - \frac{4}{p} - s_2$. We note that the power of the weight in the right side of (2.1) then satisfies:

$$-\frac{1}{2} - s_1 = p\alpha_2, \quad -\frac{1}{2} - s_2 = q\alpha_1.$$

We solve (1.10) via an iteration, and at this point, the arguments that are used are akin to those of [22], [27], and [35]. Setting $u_{-1} \equiv 0$ and $v_{-1} \equiv 0$, we recursively define u_j, v_j , $j \geq 0$ to solve

$$(3.2) \quad \begin{aligned} P_1 u_j &= F_p(v_{j-1}), & P_2 v_j &= F_q(u_{j-1}), \\ u_j(0, x) &= f_1(x), & v_j(0, x) &= f_2(x), \\ \partial_t u_j(0, x) &= g_1(x), & \partial_t v_j(0, x) &= g_2(x). \end{aligned}$$

We introduce the quantity

$$(3.3) \quad \begin{aligned} M_k(u, v) &= \|\psi_R Z^{\leq k} u\|_{\ell_q^{\alpha_1} L_t^q L_r^q L_\omega^2} + \|\psi_R Z^{\leq k} v\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2} \\ &\quad + \|\partial^{\leq k}(u, v)\|_{\ell_\infty^{-3/2} L_t^2 L_r^2 L_\omega^2} + \|\partial^{\leq k} \partial(u, v)\|_{L_t^\infty L_r^2 L_\omega^2}. \end{aligned}$$

Our first goal is to inductively show that $M_2(u_j, v_j) \leq 4C_2\varepsilon$ for some uniform constant C_2 . We first note that from (1.11) (and the assumption that the data are compactly supported) we easily obtain

$$\|\psi_R Y^{\leq 1} P_1 u_j(0, \cdot)\|_{\dot{H}^{s_1-1}} + \|\psi_R Y^{\leq 1} P_2 v_j(0, \cdot)\|_{\dot{H}^{s_2-1}} \leq C\varepsilon.$$

And thus, for C_2 chosen large enough, by (2.1),

$$(3.4) \quad \begin{aligned} & \|\psi_R Z^{\leq 2} u_j\|_{\ell_q^{\alpha_1} L_t^q L_r^q L_\omega^2} + \|\psi_R Z^{\leq 2} v_j\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2} \leq C_2\varepsilon \\ & + C\|\psi_R^p Z^{\leq 2} F_p(v_{j-1})\|_{\ell_1^{p\alpha_2} L_t^1 L_r^1 L_\omega^2} + C\|\psi_R^q Z^{\leq 2} F_q(u_{j-1})\|_{\ell_1^{q\alpha_1} L_t^1 L_r^1 L_\omega^2} \\ & + C\|\partial^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_r^2 L_\omega^2} + C\|\partial^{\leq 2} F_q(u_{j-1})\|_{L_t^1 L_r^2 L_\omega^2}. \end{aligned}$$

And from (1.11) and (1.9), we have

$$(3.5) \quad \begin{aligned} & \|\partial^{\leq 2}(u_j, v_j)\|_{\ell^{-3/2} L_t^2 L_r^2 L_\omega^2} + \|\partial^{\leq 2} \partial(u_j, v_j)\|_{L_t^\infty L_r^2 L_\omega^2} \leq C_2\varepsilon \\ & + C\|\partial^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_r^2 L_\omega^2} + C\|\partial^{\leq 2} F_q(u_{j-1})\|_{L_t^1 L_r^2 L_\omega^2}. \end{aligned}$$

From these, we first notice that $M_2(u_0, v_0) \leq 2C_2\varepsilon$, which provides the base case for the induction.

We then assume that $M_2(u_{j-1}, v_{j-1}) \leq 4C_2\varepsilon$ and show that $M_2(u_j, v_j) \leq 4C_2\varepsilon$. We first notice that (1.1) gives

$$(3.6) \quad |Z^{\leq 2} F_p(u)| \lesssim |u|^{p-1} |Z^{\leq 2} u| + |u|^{p-2} |Z^{\leq 1} u|^2.$$

By the Sobolev embeddings $H_\omega^2 \subset L_\omega^\infty$ and $H_\omega^1 \subset L_\omega^4$ on \mathbb{S}^2 and Hölder's inequality, this gives

$$\begin{aligned} \|Z^{\leq 2} F_p(v_{j-1})\|_{L_\omega^2} & \lesssim \|v_{j-1}\|_{L_\omega^\infty}^{p-1} \|Z^{\leq 2} v_{j-1}\|_{L_\omega^2} + \|v_{j-1}\|_{L_\omega^\infty}^{p-2} \|Z^{\leq 1} v_{j-1}\|_{L_\omega^4}^2 \\ & \lesssim \|Z^{\leq 2} v_{j-1}\|_{L_\omega^2}^p. \end{aligned}$$

It then immediately follows that

$$(3.7) \quad \|\psi_R^p Z^{\leq 2} F_p(v_{j-1})\|_{\ell_1^{p\alpha_2} L_t^1 L_r^1 L_\omega^2} \lesssim \|\psi_R Z^{\leq 2} v_{j-1}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2}^p \lesssim (M_2(u_{j-1}, v_{j-1}))^p \lesssim \varepsilon^p.$$

The last inequality results from the inductive hypothesis. A similar argument shows that

$$(3.8) \quad \|\psi_R^q Z^{\leq 2} F_q(u_{j-1})\|_{\ell_1^{q\alpha_1} L_t^1 L_r^1 L_\omega^2} \lesssim \varepsilon^q.$$

To finish the proof of the boundedness of $M_2(u_j, v_j)$, it remains to examine the $L_t^1 L_r^2 L_\omega^2$ pieces in the right sides of (3.4) and (3.5). The analyses will be done separately outside of a ball of radius $2R+1$, where the resulting terms will be compared to the weighted Strichartz portions of $M_2(u_{j-1}, v_{j-1})$, and inside the remaining compact set where the localized energy portions will be used.

We begin with the former using the weighted Sobolev inequalities of Lemma 2.2. Starting again at (3.6), we have

$$(3.9) \quad \begin{aligned} & \|Z^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_{r \geq 2R+1}^2 L_\omega^2} \lesssim \|r^{-\frac{\alpha_2}{p-1}} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^{\frac{2p(p-1)}{p-2}} L_\omega^\infty}^{p-1} \|r^{\alpha_2} \psi_R Z^{\leq 2} v_{j-1}\|_{L_t^p L_r^p L_\omega^2} \\ & + \left\| r^{\frac{2}{p-2}(-\alpha_2 - \frac{2}{p} + \frac{1}{2})} v_{j-1} \right\|_{L_t^p L_{r \geq 2R+1}^\infty L_\omega^\infty}^{p-2} \|r^{\alpha_2 + \frac{2}{p} - \frac{1}{2}} Z^{\leq 1} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^4 L_\omega^4}^2. \end{aligned}$$

Since (3.1) guarantees that $s_1 < 1/2$, it follows that $p\alpha_2 = -\frac{1}{2} - s_1 \geq -1$, which is equivalent to

$$-\frac{\alpha_2}{p-1} - \frac{2}{p} + \frac{p-2}{p(p-1)} \leq \alpha_2.$$

Thus, (2.3) yields

$$\|r^{-\frac{\alpha_2}{p-1}} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^{\frac{2p(p-1)}{p-2}} L_\omega^\infty} \lesssim \|r^{\alpha_2} Z^{\leq 2} v_{j-1}\|_{L_t^p L_{r \geq 2R}^p L_\omega^2}.$$

The estimate (2.4) directly applies to yield

$$\|r^{\alpha_2 + \frac{2}{p} - \frac{1}{2}} Z^{\leq 1} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^4 L_\omega^4} \lesssim \|r^{\alpha_2} Z^{\leq 2} v_{j-1}\|_{L_t^p L_{r \geq 2R}^p L_\omega^2}.$$

As above, $p\alpha_2 \geq -1$, which implies that

$$\frac{2}{p-2} \left(-\alpha_2 - \frac{2}{p} + \frac{1}{2} \right) - \frac{2}{p} \leq \alpha_2.$$

Hence (2.3) gives

$$\|r^{\frac{2}{p-2}(-\alpha_2 - \frac{2}{p} + \frac{1}{2})} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^\infty L_\omega^\infty} \lesssim \|r^{\alpha_2} Z^{\leq 2} v_{j-1}\|_{L_t^p L_{r \geq 2R}^p L_\omega^2}.$$

Plugging each of these bounds into (3.9) and using that ψ_R is identically 1 on $r \geq 2R$, it follows that

$$(3.10) \quad \|Z^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_{r \geq 2R+1}^2 L_\omega^2} \lesssim \|\psi_R Z^{\leq 2} v_{j-1}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2}^p \lesssim (M_2(u_{j-1}, v_{j-1}))^p \lesssim \varepsilon^p.$$

And an analogous argument in the symmetric variable q shows that

$$(3.11) \quad \|Z^{\leq 2} F_q(u_{j-1})\|_{L_t^1 L_{r \geq 2R+1}^2 L_\omega^2} \lesssim \varepsilon^q,$$

which leaves the analysis of the $L_t^1 L_r^2 L_\omega^2$ pieces over the region $r \leq 2R+1$ where the coefficients of Z are bounded.

We again start with (3.6) and apply Hölder's inequality to obtain

$$(3.12) \quad \begin{aligned} \|Z^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_{r \leq 2R+1}^2 L_\omega^2} &\lesssim \|v_{j-1}\|_{L_t^\infty L_r^\infty L_\omega^\infty}^{p-2} \|v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^\infty L_\omega^\infty} \|\partial^{\leq 2} v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^2 L_\omega^2} \\ &\quad + \|v_{j-1}\|_{L_t^\infty L_r^\infty L_\omega^\infty}^{p-2} \|\partial^{\leq 1} v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^4 L_\omega^4}^2. \end{aligned}$$

By Sobolev embeddings, we have

$$\|v_{j-1}\|_{L_t^\infty L_r^\infty L_\omega^\infty} \lesssim \|\partial^{\leq 1} v_{j-1}\|_{L_t^\infty L_r^6 L_\omega^6} \lesssim \|\partial^{\leq 1} \partial v_{j-1}\|_{L_t^\infty L_r^2 L_\omega^2}.$$

Similarly, Sobolev embeddings (with a localizing factor) give

$$\|v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^\infty L_\omega^\infty} \lesssim \|\partial^{\leq 2} v_{j-1}\|_{L_t^2 L_{r \leq 2R+2}^2 L_\omega^2} \lesssim \|\partial^{\leq 2} v_{j-1}\|_{\ell_\infty^{-3/2} L_t^2 L_r^2 L_\omega^2}$$

and

$$\|\partial^{\leq 1} v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^4 L_\omega^4} \lesssim \|\partial^{\leq 2} v_{j-1}\|_{L_t^2 L_{r \leq 2R+2}^2 L_\omega^2} \lesssim \|\partial^{\leq 2} v_{j-1}\|_{\ell_\infty^{-3/2} L_t^2 L_r^2 L_\omega^2}.$$

These bounds in (3.12) show that

$$(3.13) \quad \begin{aligned} \|Z^{\leq 2} F_p(v_{j-1})\|_{L_t^1 L_{r \leq 2R+1}^2 L_\omega^2} &\lesssim \|\partial^{\leq 1} \partial v_{j-1}\|_{L_t^\infty L_r^2 L_\omega^2}^{p-2} \|\partial^{\leq 2} v_{j-1}\|_{\ell_\infty^{-3/2} L_t^2 L_r^2 L_\omega^2}^2 \\ &\lesssim (M_2(u_{j-1}, v_{j-1}))^p \lesssim \varepsilon^p. \end{aligned}$$

And an analogous argument gives

$$(3.14) \quad \|Z^{\leq 2} F_q(u_{j-1})\|_{L_t^1 L_{r \leq 2R+1}^2 L_\omega^2} \lesssim \varepsilon^q.$$

Using (3.7), (3.8), (3.10), (3.11), (3.13), and (3.14) in (3.4) and (3.5), we obtain

$$M_2(u_j, v_j) \leq 2C_2\varepsilon + C_3\varepsilon^{\min(p,q)}$$

for some constant C_3 that is independent of j . Since $p, q \geq 2$, if ε is sufficiently small, the desired bound $M_2(u_j, v_j) \leq 4C_2\varepsilon$ follows.

It remains to show that the sequence converges. To do so, we shall show

$$(3.15) \quad M_0(u_j - u_{j-1}, v_j - v_{j-1}) \leq \frac{1}{2} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}),$$

and this will complete the proof. Applying (2.2) and (1.9), we have

$$(3.16) \quad \begin{aligned} & \|\psi_R(u_j - u_{j-1})\|_{\ell_1^{\alpha_1} L_t^q L_r^q L_\omega^2} + \|\psi_R(v_j - v_{j-1})\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2} \\ & \lesssim \|\psi_R^p(F_p(v_{j-1}) - F_p(v_{j-2}))\|_{\ell_1^{p\alpha_2} L_t^1 L_r^1 L_\omega^2} + \|\psi_R^q(F_q(u_{j-1}) - F_q(u_{j-2}))\|_{\ell_1^{q\alpha_1} L_t^1 L_r^1 L_\omega^2} \\ & \quad + \|F_p(v_{j-1}) - F_p(v_{j-2})\|_{L_t^1 L_r^2 L_\omega^2} + \|F_q(u_{j-1}) - F_q(u_{j-2})\|_{L_t^1 L_r^2 L_\omega^2}, \end{aligned}$$

and

$$(3.17) \quad \begin{aligned} & \|(u_j - u_{j-1}, v_j - v_{j-1})\|_{\ell^{-3/2} L_t^2 L_r^2 L_\omega^2} + \|\partial(u_j - u_{j-1}, v_j - v_{j-1})\|_{L_t^\infty L_r^2 L_\omega^2} \\ & \lesssim \|F_p(v_{j-1}) - F_p(v_{j-2})\|_{L_t^1 L_r^2 L_\omega^2} + \|F_q(u_{j-1}) - F_q(u_{j-2})\|_{L_t^1 L_r^2 L_\omega^2}. \end{aligned}$$

Noting that

$$(3.18) \quad |F_p(v_{j-1}) - F_p(v_{j-2})| \lesssim (|v_{j-1}|^{p-1} + |v_{j-2}|^{p-1})|v_{j-1} - v_{j-2}|,$$

we can quickly observe that

$$(3.19) \quad \begin{aligned} & \|\psi_R^p(F_p(v_{j-1}) - F_p(v_{j-2}))\|_{\ell_1^{p\alpha_2} L_t^1 L_r^1 L_\omega^2} \\ & \lesssim \left(\|\psi_R v_{j-1}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^\infty}^{p-1} + \|\psi_R v_{j-2}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^\infty}^{p-1} \right) \|\psi_R(v_{j-1} - v_{j-2})\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2} \\ & \lesssim \left(\|\psi_R Z^{\leq 2} v_{j-1}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2}^{p-1} + \|\psi_R Z^{\leq 2} v_{j-2}\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2}^{p-1} \right) \\ & \quad \times \|\psi_R(v_{j-1} - v_{j-2})\|_{\ell_p^{\alpha_2} L_t^p L_r^p L_\omega^2} \\ & \lesssim (4C_2\varepsilon)^{p-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}). \end{aligned}$$

A similar argument yields

$$(3.20) \quad \|\psi_R^q(F_q(u_{j-1}) - F_q(u_{j-2}))\|_{\ell_1^{q\alpha_1} L_t^1 L_r^1 L_\omega^2} \lesssim (4C_2\varepsilon)^{q-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}).$$

We also start at (3.18) to control the $L_t^1 L_r^2 L_\omega^2$ terms. There we get, using (2.3) as above,

$$(3.21) \quad \begin{aligned} & \|F_p(v_{j-1}) - F_p(v_{j-2})\|_{L_t^1 L_{r \geq 2R+1}^2 L_\omega^2} \\ & \lesssim \left(\|r^{-\frac{\alpha_2}{p-1}} v_{j-1}\|_{L_t^p L_{r \geq 2R+1}^{\frac{2p(p-1)}{p-2}} L_\omega^\infty}^{p-1} + \|r^{-\frac{\alpha_2}{p-1}} v_{j-2}\|_{L_t^p L_{r \geq 2R+1}^{\frac{2p(p-1)}{p-2}} L_\omega^\infty}^{p-1} \right) \|r^{\alpha_2} \psi_R(v_{j-1} - v_{j-2})\|_{L_t^p L_r^p L_\omega^2} \\ & \lesssim [(M_2(u_{j-1}, v_{j-1}))^{p-1} + (M_2(u_{j-2}, v_{j-2}))^{p-1}] M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}) \\ & \lesssim \varepsilon^{p-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}). \end{aligned}$$

And similarly

$$(3.22) \quad \|F_q(u_{j-1}) - F_q(u_{j-2})\|_{L_t^1 L_{r \geq 2R+1}^2 L_\omega^2} \lesssim \varepsilon^{q-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}).$$

Finally, using Sobolev embeddings as in (3.13),

$$(3.23) \quad \begin{aligned} & \|F_p(v_{j-1}) - F_p(v_{j-2})\|_{L_t^1 L_{r \leq 2R+1}^2 L_\omega^2} \\ & \lesssim [\|v_{j-1}\|_{L_t^\infty L_r^\infty L_\omega^\infty}^{p-2} \|v_{j-1}\|_{L_t^2 L_{r \leq 2R+1}^\infty L_\omega^\infty} + \|v_{j-2}\|_{L_t^\infty L_r^\infty L_\omega^\infty}^{p-2} \|v_{j-2}\|_{L_t^2 L_{r \leq 2R+1}^\infty L_\omega^\infty}] \\ & \quad \times \|v_{j-1} - v_{j-2}\|_{L_t^2 L_{r \leq 2R+1}^2 L_\omega^2} \\ & \lesssim [(M_2(u_{j-1}, v_{j-1}))^{p-1} + (M_2(u_{j-2}, v_{j-2}))^{p-1}] M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}) \\ & \lesssim \varepsilon^{p-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}), \end{aligned}$$

and by the same procedures

$$(3.24) \quad \|F_q(u_{j-1}) - F_q(u_{j-2})\|_{L_t^1 L_{r \leq 2R+1}^2 L_\omega^2} \lesssim \varepsilon^{q-1} M_0(u_{j-1} - u_{j-2}, v_{j-1} - v_{j-2}).$$

Plugging (3.19)-(3.24) into (3.16) and (3.17) immediately yields (3.15) provided that ε is sufficiently small, and this completes the proof.

REFERENCES

- [1] Rentaro Agemi, Yuki Kurokawa, and Hiroyuki Takamura. Critical curve for p - q systems of nonlinear wave equations in three space dimensions. *J. Differential Equations*, 167(1):87–133, 2000.
- [2] Serge Alinhac. On the Morawetz–Keel–Smith–Sogge inequality for the wave equation on a curved background. *Publ. Res. Inst. Math. Sci.*, 42(3):705–720, 2006.
- [3] Lars Andersson and Pieter Blue. Hidden symmetries and decay for the wave equation on the Kerr spacetime. *Ann. of Math. (2)*, 182(3):787–853, 2015.
- [4] P. Blue and A. Soffer. Errata for “Global existence and scattering for the nonlinear Schrödinger equation on Schwarzschild manifolds”, “Semilinear wave equations on the Schwarzschild manifold I: Local Decay Estimates”, and “The wave equation on the Schwarzschild metric II: Local decay for the spin 2 Regge Wheeler equation. Preprint. ArXiv:gr-qc/0608073.
- [5] P. Blue and A. Soffer. Semilinear wave equations on the Schwarzschild manifold. I. Local decay estimates. *Adv. Differential Equations*, 8(5):595–614, 2003.
- [6] Jean-François Bony and Dietrich Häfner. The semilinear wave equation on asymptotically Euclidean manifolds. *Comm. Partial Differential Equations*, 35(1):23–67, 2010.
- [7] Robert Booth, Hans Christianson, Jason Metcalfe, and Jacob Perry. Localized energy for wave equations with degenerate trapping. In preparation.
- [8] N. Burq. Global Strichartz estimates for nontrapping geometries: about an article by H. F. Smith and C. D. Sogge: “Global Strichartz estimates for nontrapping perturbations of the Laplacian” [Comm. Partial Differential Equation **25** (2000), no. 11-12 2171–2183; MR1789924 (2001j:35180)]. *Comm. Partial Differential Equations*, 28(9-10):1675–1683, 2003.
- [9] M. Dafermos and I. Rodnianski. Decay for solutions of the wave equation on Kerr exterior spacetimes I-II: The cases $|a| \ll M$ or axisymmetry. Preprint. ArXiv:1010.5132.
- [10] M. Dafermos and I. Rodnianski. A note on energy currents and decay for the wave equation on a Schwarzschild background. Preprint. ArXiv:0710.0171.
- [11] Mihalis Dafermos and Igor Rodnianski. The red-shift effect and radiation decay on black hole spacetimes. *Comm. Pure Appl. Math.*, 62(7):859–919, 2009.
- [12] Mihalis Dafermos and Igor Rodnianski. A proof of the uniform boundedness of solutions to the wave equation on slowly rotating Kerr backgrounds. *Invent. Math.*, 185(3):467–559, 2011.
- [13] Mihalis Dafermos, Igor Rodnianski, and Yakov Shlapentokh-Rothman. Decay for solutions of the wave equation on Kerr exterior spacetimes III: The full subextremal case $|a| < M$. *Ann. of Math. (2)*, 183(3):787–913, 2016.
- [14] D. Del Santo. Global existence and blow-up for a hyperbolic system in three space dimensions. *Rend. Istit. Mat. Univ. Trieste*, 29(1-2):115–140 (1998), 1997.

- [15] D. Del Santo and È. Mitidieri. Blow-up of solutions of a hyperbolic system: the critical case. *Differ. Uravn.*, 34(no. 9):1155–1161, 1293, 1998.
- [16] Daniele Del Santo, Vladimir Georgiev, and Enzo Mitidieri. Global existence of the solutions and formation of singularities for a class of hyperbolic systems. In *Geometrical optics and related topics (Cortona, 1996)*, volume 32 of *Progr. Nonlinear Differential Equations Appl.*, pages 117–140. Birkhäuser Boston, Boston, MA, 1997.
- [17] Keng Deng. Nonexistence of global solutions of a nonlinear hyperbolic system. *Trans. Amer. Math. Soc.*, 349(4):1685–1696, 1997.
- [18] Yi Du, Jason Metcalfe, Christopher D. Sogge, and Yi Zhou. Concerning the Strauss conjecture and almost global existence for nonlinear Dirichlet-wave equations in 4-dimensions. *Comm. Partial Differential Equations*, 33(7-9):1487–1506, 2008.
- [19] Daoyuan Fang and Chengbo Wang. Weighted Strichartz estimates with angular regularity and their applications. *Forum Math.*, 23(1):181–205, 2011.
- [20] Vladimir Georgiev, Hans Lindblad, and Christopher D. Sogge. Weighted Strichartz estimates and global existence for semilinear wave equations. *Amer. J. Math.*, 119(6):1291–1319, 1997.
- [21] Vladimir Georgiev, Hiroyuki Takamura, and Zhou Yi. The lifespan of solutions to nonlinear systems of a high-dimensional wave equation. *Nonlinear Anal.*, 64(10):2215–2250, 2006.
- [22] Kunio Hidano, Jason Metcalfe, Hart F. Smith, Christopher D. Sogge, and Yi Zhou. On abstract Strichartz estimates and the Strauss conjecture for nontrapping obstacles. *Trans. Amer. Math. Soc.*, 362(5):2789–2809, 2010.
- [23] Fritz John. Blow-up of solutions of nonlinear wave equations in three space dimensions. *Manuscripta Math.*, 28(1-3):235–268, 1979.
- [24] S. Klainerman. The null condition and global existence to nonlinear wave equations. In *Nonlinear systems of partial differential equations in applied mathematics, Part 1 (Santa Fe, N.M., 1984)*, volume 23 of *Lectures in Appl. Math.*, pages 293–326. Amer. Math. Soc., Providence, RI, 1986.
- [25] Hideo Kubo and Masahito Ohta. Critical blowup for systems of semilinear wave equations in low space dimensions. *J. Math. Anal. Appl.*, 240(2):340–360, 1999.
- [26] Hideo Kubo and Masahito Ohta. On the global behavior of classical solutions to coupled systems of semilinear wave equations. In *New trends in the theory of hyperbolic equations*, volume 159 of *Oper. Theory Adv. Appl.*, pages 113–211. Birkhäuser, Basel, 2005.
- [27] Hans Lindblad, Jason Metcalfe, Christopher D. Sogge, Mihai Tohaneanu, and Chengbo Wang. The Strauss conjecture on Kerr black hole backgrounds. *Math. Ann.*, 359(3-4):637–661, 2014.
- [28] Jeremy Marzuola, Jason Metcalfe, Daniel Tataru, and Mihai Tohaneanu. Strichartz estimates on Schwarzschild black hole backgrounds. *Comm. Math. Phys.*, 293(1):37–83, 2010.
- [29] Jason Metcalfe and Christopher D. Sogge. Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods. *SIAM J. Math. Anal.*, 38(1):188–209, 2006.
- [30] Jason Metcalfe and Christopher D. Sogge. Global existence of null-form wave equations in exterior domains. *Math. Z.*, 256(3):521–549, 2007.
- [31] Jason Metcalfe, Jacob Sterbenz, and Daniel Tataru. Local energy decay for scalar fields on time dependent non-trapping backgrounds. In preparation.
- [32] Jason Metcalfe and Daniel Tataru. Decay estimates for variable coefficient wave equations in exterior domains. In *Advances in phase space analysis of partial differential equations*, volume 78 of *Progr. Nonlinear Differential Equations Appl.*, pages 201–216. Birkhäuser Boston, Inc., Boston, MA, 2009.
- [33] Jason Metcalfe and Daniel Tataru. Global parametrices and dispersive estimates for variable coefficient wave equations. *Math. Ann.*, 353(4):1183–1237, 2012.
- [34] Jason Metcalfe, Daniel Tataru, and Mihai Tohaneanu. Price’s law on nonstationary space-times. *Adv. Math.*, 230(3):995–1028, 2012.
- [35] Jason Metcalfe and Chengbo Wang. The Strauss conjecture on asymptotically flat space-times. *arXiv preprint arXiv:1605.02157*, 2016.
- [36] Cathleen S. Morawetz. Time decay for the nonlinear Klein-Gordon equations. *Proc. Roy. Soc. Ser. A*, 306:291–296, 1968.
- [37] James V. Ralston. Solutions of the wave equation with localized energy. *Comm. Pure Appl. Math.*, 22:807–823, 1969.
- [38] Jan Sbierski. Characterisation of the energy of Gaussian beams on Lorentzian manifolds: with applications to black hole spacetimes. *Anal. PDE*, 8(6):1379–1420, 2015.

- [39] Jack Schaeffer. The equation $u_{tt} - \Delta u = |u|^p$ for the critical value of p . *Proc. Roy. Soc. Edinburgh Sect. A*, 101(1-2):31–44, 1985.
- [40] Thomas C. Sideris. Nonexistence of global solutions to semilinear wave equations in high dimensions. *J. Differential Equations*, 52(3):378–406, 1984.
- [41] Hart F. Smith, Christopher D. Sogge, and Chengbo Wang. Strichartz estimates for Dirichlet-wave equations in two dimensions with applications. *Trans. Amer. Math. Soc.*, 364(6):3329–3347, 2012.
- [42] Christopher D. Sogge and Chengbo Wang. Concerning the wave equation on asymptotically Euclidean manifolds. *J. Anal. Math.*, 112:1–32, 2010.
- [43] Jacob Sterbenz. Angular regularity and Strichartz estimates for the wave equation. *Int. Math. Res. Not.*, (4):187–231, 2005. With an appendix by Igor Rodnianski.
- [44] Walter A. Strauss. Nonlinear scattering theory at low energy. *J. Funct. Anal.*, 41(1):110–133, 1981.
- [45] Daniel Tataru. Strichartz estimates in the hyperbolic space and global existence for the semilinear wave equation. *Trans. Amer. Math. Soc.*, 353(2):795–807 (electronic), 2001.
- [46] Daniel Tataru. Local decay of waves on asymptotically flat stationary space-times. *Amer. J. Math.*, 135(2):361–401, 2013.
- [47] Daniel Tataru and Mihai Tohaneanu. A local energy estimate on Kerr black hole backgrounds. *Int. Math. Res. Not. IMRN*, (2):248–292, 2011.
- [48] Chengbo Wang. Long time existence for semilinear wave equations on asymptotically flat space-times. *arXiv preprint arXiv:1504.05652*, 2015.
- [49] Chengbo Wang and Xin Yu. Concerning the Strauss conjecture on asymptotically Euclidean manifolds. *J. Math. Anal. Appl.*, 379(2):549–566, 2011.
- [50] Chengbo Wang and Xin Yu. Recent works on the Strauss conjecture. In *Recent advances in harmonic analysis and partial differential equations*, volume 581 of *Contemp. Math.*, pages 235–256. Amer. Math. Soc., Providence, RI, 2012.
- [51] Borislav T. Yordanov and Qi S. Zhang. Finite time blow up for critical wave equations in high dimensions. *J. Funct. Anal.*, 231(2):361–374, 2006.
- [52] Xin Yu. Generalized Strichartz estimates on perturbed wave equation and applications on Strauss conjecture. *Differential Integral Equations*, 24(5-6):443–468, 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27599-3250, USA

E-mail address: metcalfe@email.unc.edu

URL: <http://metcalfe.web.unc.edu>

UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27599, USA

E-mail address: davidspencer6174@gmail.com